

WARNINGS:

WITHHOLDING PERIOD: Number of days which should be allowed between the last application and harvesting or grazing (feeding) of the crop:
Apples, Pears and Olives..... 14 days

- Handle with care.
- Irritating to the eyes and mildly irritating to the skin.
- Handle with care.
- May cause eye irritation and skin sensitization.
- Harmful if inhaled or swallowed.
- Toxic to fish and wildlife.
- Store away from food and feeds.
- Store in the original container/bag, tightly closed, in a cool, dry place.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Prevent contamination of food, feed, drinking water and eating utensils.
- Do not eat, drink or smoke whilst handling or applying, or before washing hands and face and changing clothing.
- Store away from food and feed, and keep out of reach of children and animals Safety during mixing:

Safety during mixing:

- Wear eye protection (shield or spectacles). In the event of eye contamination, flush out immediately.
- Wear protective rubber gloves. If you get it on your skin, wash it off immediately.
- Wear protective clothing. In the event of spillage on clothes, undress and wash.
- Rinse empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the spray tank. Destroy or suitably dispose the container. Never reuse an empty container for any other purpose.

Safety during application:

- Avoid contact of spray with skin or eyes.
- Do not breathe spray mist.
- Prevent spreading of spray mist to other crops, pastures, rivers, dams and areas which are not under treatment.

Safety after application:

- Clean applicator thoroughly after use and dispose of wash water where it will not contaminate other crops, pastures, rivers or dams.
- Destroy empty container and do not use for any other purpose.
- Remove work clothes, wash yourself and wash clothes thoroughly.



FARMAG CAPTAN 500 SC

Reg No L 10311 Act / Wet 36 of 1947

A suspension concentrate fungicide for the control of pests in crops as indicated.

'n Suspensiekonsentraat swamdoder vir die beheer van plae in gewasse soos aangedui.

FRAC FUNGICIDE GROUP CODE:	M4	FRAC SWAMDODER GROEPKODE:
----------------------------	----	---------------------------

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Captab (dicarboximide)..... 500 g/l..... Captab (dikarboksimide)

Manufacture Date:
Vervaardigingsdatum:

Batch No.:
Lotnommer:

NET MASS/ INHOUD

**SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBLIJET VIR VOLLE BESONDERHEDE**

REGISTRATION HOLDER/REGISTRASIEHOUER:

Cedar Falls Properties 130 (Pty) Ltd.

Co. Reg.No. 2006/031383/07

22 Burnside Drive, Old Mill Industrial Park, Mount Edgecombe, South Africa, 4300 • Tel. (031) 502 4196

• Emergency Number: Griffon Poison Information Centre

UN No.: 3082



CAUTION
VERSIGTIG



165 x 140

CYAN

MAG

YEL

BLK

DIRECTIONS FOR USE: Use only as indicated

CROP / DISEASE	DOSAGE	REMARKS
APPLES AND PEARS Scab (<i>Venturia inaequalis</i>)	150 - 200ml / 100 l water	<p>Pre-blossom: Start application at green-tip stage and repeat at 7 day intervals.</p> <p>Post-blossom: Continue with 3 - 4 applications depending on weather conditions and with 10 - 14 day intervals. Use the higher dosage rate under high infestation pressure.</p> <p>Note: FARMAG CAPTAN 500 SC may cause leafspots on Pearmain, Delicious, Cox and other sensitive cultivars</p>
OLIVES Leaf spot (<i>Cyclococonium oleaginum</i>)	150 - 200ml / 100l water	Apply two high volume applications to the point of runoff. Apply one in spring before the new flush appears and then a second in autumn before the start of winter.
TOBACCO SEEDBEDS Anthracnose (<i>Colletotrichum destructivum</i>)		Apply 15 l / 10m ² as a soil drench 3 weeks after germination and wash into the soil with 40 l of water.
Black root rot (<i>Thielaviopsis vasicola</i>)		Repeat applications every 10 – 14 days until transplanting.
Black shank (<i>Phytophthora nicotianae</i>)		
Damping-off (<i>Pythium, Fudarium and Rhizoctonia spp</i>)	200ml / 100l water	

WAARSKUWINGS:

ONTHOUDEINGS PERIODE: Aantal dae wat toegelaat behoort te word tussen die laaste toediening en oes of beweiding (voeding) van die gewas:
Appels, Pere en Olywe.....14 dae

- Hanteer versigtig.
- Mag oog irritasie en velgevoeligheid veroorsaak.
- Nadelig wanneer ingeasem of ingesluk word.
- Giftig vir vise en wildlewe.
- Berg weg van voedsel en voer.
- Berg in die oorspronklike houer/sak, dig toegemaak, in 'n koel, droë plek.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit doeltreffend sal wees onder alle toestande nie aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormalle klimaats- en beringstoestande; kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstsnd van die siekte teen die betrokke middel sowel as die metode, tyd en akkuraathed van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroeи, die omgewing of vir nadelige effek op mens of dier of vir gebrek aan prestatie van die betrokke middel as gevolg van die versuim van die gebruiker om etiket aanwysings te volg of as gevolg van die ontstaan van toestande wat nie voorsien kon word in die registrasie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMAATREËLS:

- Verhoed besoedeling van voedsel, voer, drinkwater en eetgerei.
- Moenie eet, drink of rook tydens hantering of toediening, of voordat hande en gesig gewas is en klere verwissel is.
- Berg weg van voedsel en voer en hou buite bereik van kinders en diere,

Veiligheid tydens vermenging:

- Dra oog beskerming (skild of bril). In geval van oog kontaminasie, spoel onmiddellik uit.
- Dra beskermende rubber handskoene. In geval van velkontak, was dit onmiddellik af.
- Dra beskermende klere. In geval van storting op klere, trek uit en was.
- Spoel leë houer drie maal met 'n volume gelykstaande tot 'n minimum van 10% van die houer. Voeg die spoelings water by die inhoud van die spuitenk. Vernietig die houer. Moet nooit die leë houer vir enige ander doel hergebruik nie,

Veiligheid tydens toediening:

- Vermy kontak van spuitmis met vel of oë.
- Moenie spuitmis inasem nie.
- Vermy verspreiding van sputnewel na ander gewasse, weidings, riviere, damme en areas nie onder behandeling nie.

Veiligheid na toediening:

- Maak toedieningsapparaat deeglik skoon na gebruik en doen weg met waswater waar dit nie ander gewasse, weidings, riviere of damme sal besoeedel nie.
- Vernietig leë houer en moet dit vir geen ander doel gebruik nie.
- Trek werksklere uit, was jouself en was klere deeglik.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui

GEWAS / SIEKTE	DOSIS	OPMERKINGS
APPELS AND PERE Skurf (<i>Venturia inaequalis</i>)	150 - 200mℓ / 100ℓ water	<p>Pre-bloeisel: Begin toediening by groenpunt stadium en herhaal met 7 dae intervalle.</p> <p>Post-bloeisel: Gaan voort met 3 - 4 bespuitings afhangende van die weersomstandighede en met 10 - 14 dag intervalle. Gebruik die hoër dosis hoë infestasie druk.</p> <p>Nota: FARMAG CAPTAN 500 SC mag blaarvlek op Pearmain, Delicious, Cox en ander sensitiewe appel kultivars veroorsaak.</p>
OLYWE Blaarvlek (<i>Cycloconium oleaginum</i>)	150 - 200mℓ / 100ℓ water	Dien twee hoë volume bespuitings tot die punt van afloop toe. Dien een in die lente voor die nuwe groei plaasvind en dan 'n tweede in herfs voor aanvang van die winter.
TABAK SAADBED Anthraknose (<i>Colletotrichum destructivum</i>)		Dien 15 ℥ / 10m ² as 'n gronddeurweking 3 weke na ontkieming toe en was in die grond in met 40 ℥ water.
Swart wortel vrot (<i>Theilaviopsis vasicola</i>)		
Swart stam (<i>Phytophthora nicotianae</i>)	200mℓ / 100ℓ water	Herhaal toedienings elke 10 – 14 dae tot en met uitplanting.
Verwelksiekte (<i>Pythium, Fudarium en Rhizoctonia spp</i>)		



FARMAG CAPTAN 500 SC

Reg No L 10311 Act / Wet 36 of 1947

A suspension concentrate fungicide for the control of pests in crops as indicated.

'n Suspensiekonsentraat swamdoder vir die beheer van plae in gewasse soos aangedui.

FRAC FUNGICIDE GROUP CODE:	M4	FRAC SWAMDODER GROEPKODE:
----------------------------	----	---------------------------

ACTIVE INGREDIENT/AKTIEWE BESTANDDEEL:

Captab (dicarboximide)..... 500 g/l..... Captab (dikarboksimiede)

Manufacture Date:
Vervaardigingsdatum:

Batch No.:
Lotnommer:

NET MASS/ INHOUD

**SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBLIJET VIR VOLLE BESONDERHEDE**

REGISTRATION HOLDER/REGISTRASIEHOUER:

Cedar Falls Properties 130 (Pty) Ltd.

Co. Reg.No. 2006/031383/07

22 Burnside Drive, Old Mill Industrial Park, Mount Edgecombe, South Africa, 4300 • Tel. (031) 502 4196

• Emergency Number: Griffon Poison Information Centre

UN No.: 3082



CAUTION
VERSIGTIG

