



MONO POTASSIUM PHOSPHATE

Reg. No.: K 9954 Act /Wet No. 36 of/van 1947

A water soluble plant nutrient

N' Wateroplosbare plantvoedingstof

FERTILIZER GROUP	2	MISSTOF GROEP
------------------	---	---------------

PLANT NUTRIENT CONTENT / PLANTVOEDINGSSTOFINHOUD:

Potassium.....282 g/kg..... kalium
Phosphate.....227 g/kg..... fosfaat

Manufacture date:
Vervaardigingsdatum:

Batch No.:
Lotnommer:

CONTENTS/INHOUD

**SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE**

REGISTRATION HOLDER/ REGISTRASIEHOUER

Cedar Fall Properties 130 (Pty) Ltd
Co. Reg. No: 2006/031383/07
22 Burnside Drive
Old Mill Industrial Park
South Africa 4300
Tel: (+27) 031 5388700

DIRECTIONS OF USE:

MONO POTASSIUM PHOSPHATE is recommended for use during plant growth periods that require relatively high potassium and phosphorus levels, such as during flowering and fruiting periods, and during periods when the uptake and movement of potassium and phosphorus in the plant is reduced. This product is specifically formulated to correct potassium and phosphorus deficiencies. This product is a completely water soluble fertiliser and chlorine free. This product is readily taken up by the plant and is formulated for use as a foliar feed, for fertigation, and side dressing. **MONO POTASSIUM PHOSPHATE** has a very low conductivity value and is suited for use in irrigation water with high salt indexes. **MONO POTASSIUM PHOSPHATE** is compatible with most agricultural remedies.

It is however advisable to do a miscibility test prior to mixing with other chemicals. **MONO POTASSIUM PHOSPHATE** should preferably be applied in the early morning or late afternoon.

Do not apply to plants that are undergoing a period of moisture stress.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED		
APPLICATION	RATE PER HECTARE	REMARKS
Foliar feed	1 – 5 kg 1 – 3 kg 5 - 10 kg	Vegetable and Ornamental crops: Apply as a full cover spray. Apply at 1 – 2 week intervals as required. Aerial Application: Apply at 1 – 2 week intervals as required. Large trees, Orchards and Vineyards: Apply at 2 – 4 week intervals. In the case of deficiencies repeat treatment one week later.
Fertigation (continuous)	50 – 100 kg	Use lower concentrations for young plants and increase the rate as the plants increase in size.
Side Dressing	50 – 100 kg	Vegetable and Ornamental crops: Apply in the row. Depending on the intensity of cultivation the application can be repeated 3 – 4 weeks later.

Note: The rate of application is dependent on the soil fertility and reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies. Consult with your chemical distributor, consultant or manufacturer should crop specific programs or any other information be required around the use of fertilizers.

GEBRUIKSAANWYSINGS:

MONO POTASSIUM PHOSPHATE word aanbeveel vir gebruik tydens plant groei periodes wat relatief hoë vlakke kalium en fosfaat vereis, soos gedurende blom en vrugdraende periodes. En tydens periodes wanneer die opname en beweging van kalium en fosfaat in die plant verminder is. Hierdie produk is spesifiek geformuleer om kalium en fosfaat tekorte reg te stel. Die produk is 'n volkome water oplosbare kunsmis en is chloor vry. Hierdie produk word gereedlik deur die plant opgeneem is geformuleer vir gebruik as 'n blaarvoeder, vir sproeibemesting en formulated for use as a foliar feed, for fertigation, en sybewerking. **MONO POTASSIUM PHOSPHATE** het 'n baie lae geleidings waarde en is geskik vir gebruik in besproeiings water met hoë sout indekse. **MONO POTASSIUM PHOSPHATE** is verenigbaar met meeste landboumiddels.

Dit word egter aanbeveel om 'n verenigbaarheids toets te doen voordat met ander chemikalieë gemeng word. MONO POTASSIUM PHOSPHATE behoort verkieslik vroeg in die môre of laat in die middag toegedien te word.

Moenie toedien op plante wat 'n periode van vogtigheid stres ondergaan nie.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS VOORGESKRYF		
TOEDIENING	DOSIS PER HEKTAAR	OPMERKINGS
Blaarvoeding	1 – 5 kg 1 – 3 kg 5 - 10 kg	Groente en Sierplant gewasse: Dien toe as 'n vol bedekte bespuiting. Dien toe met 1 – 2 week intervalle soos benodig. Lugtoediening: Dien toe met 1 – 2 week intervalle soos benodig. Groot bome, Boorde en Wingerde: Dien toe met 2 – 4 week intervalle. In die geval van tekorte herhaal behandeling een week later.
Sproeibemesting (deurlopend)	50 – 100 kg	Gebruik laer konsentrasies vir jong plante en verhoog die dosis soos die plante toeneem in grootte.
Sybewerking	50 – 100 kg	Groente en Sierplant gewasse: Dien toe in die ry. Afhangend van die intensiteit van bewerking kan die toediening 3 – 4 weke later herhaal word.

Nota: Die dosis van toediening hang af van die grond vrugbaarheid en die reaksie vereis. Die laer dosisse moet gebruik word vir onderhoud en die hoër dosisse sal voedings tekorte korrigeer. Raadpleeg jou chemiese verspreidings agent, adviseur of vervaardiger vir gewas spesifieke programme of enige ander inligting vereis rakende die gebruik van misstowwe.