

WARNINGS:

- Harmful when swallowed.
- Concentrate may cause irritation to the eyes and skin.
- **FLAMMABLE** – do not store near open flame.
- Store in a cool, dry place.
- Store away from food, feed, seed and other agricultural chemicals.
- Keep out of reach of children, uninformed persons and animals.
- May be toxic to fish.
- **Re-entry:** Do not enter treated area within 1 day after treatment unless wearing protective clothing.
- **In case of poisoning call a physician and make this label available to him/her.**

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions, because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of water, compatibility with other substances not indicated on the label and the occurrence of resistance of the weed against the remedy, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, and the environment or harm to man or animal, or for lack of performance of the remedy concerned due to failure by the user to follow the label instructions, or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

For your own safety take the following precautions:

- Do not eat, drink or smoke while handling this product.
- Do not eat, drink or smoke while mixing or applying the product or before washing hands and face and changing clothing.
- After using this product wash hands and face.
- Prevent contamination of food, feeds, drinking water and eating utensils.

When mixing:

- Wear eye protection and impermeable gloves.
- Wash with soap and running water after accidental skin contact.
- In case of eye contact, flush eyes immediately with clean running water, and seek medical advice.

When spraying:

- Do not inhale fumes or spray mist.
- Avoid contact with the spray solution.
- Wear suitable protective clothing.
- Prevent drift onto other crops, grazing, rivers, dams or areas not under treatment.

After spraying:

- Clean applicator thoroughly after use and dispose of rinsing water where it will not contaminate food, crops, grazing, rivers or dams.
- Change and wash your work clothes.
- Wash your exposed skin area and entire body, including your hair and under your finger nails with soap and water.

- Triple rinse empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow to drain for at least 30 seconds after the flow has slowed down to a drip. Thereafter, rinse the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsings to the contents of the spray tank before destroying the container in the described manner.
- Destroy the empty container by perforation and flattening and dispose of it in a safe manner.
- Never re-use the empty container for any other purpose.

SYMPTOMS OF HUMAN POISONING:

Symptoms relating to use:

Skin Contact: Irritating to skin (itching, redness, pain).

Eye Contact: Not expected to present a significant eye contact hazard under anticipated conditions of normal use.

Ingestion: Abdominal pain, nausea, breathing difficulties, harmful. May cause lung damage if swallowed.

FIRST AID TREATMENT:

Skin contact: For even minor contact, remove contaminated clothing, shoes and leather goods. Wash skin gently and thoroughly with water and non-abrasive soap. If irritation persists, obtain medical attention.

Eye contact: Immediately flush the contaminated eyes with gently flowing clean water for 20 minutes, holding the eyelid(s) open until no evidence of chemical remains. If irritation persists, obtain medical attention.

Inhalation: Assure fresh air breathing. Remove source of contamination or move victim to fresh air. If breathing is difficult, administer oxygen. If breathing has stopped, give artificial respiration. If breathing difficulty persists, obtain medical attention.

Ingestion: Do not induce vomiting. Have victim rinse mouth thoroughly with water. Obtain medical advice.

NOTE TO PHYSICIAN: No specific antidote is available. Treat symptomatically and supportively when required.

RESISTANCE WARNING:

VULTURE 480 EC is a group code O herbicide. Any weed population may contain individuals naturally resistant to **VULTURE 480 EC** and any other group code O herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by **VULTURE 480 EC** or any other group code O herbicide.

To delay herbicide resistance:

- Avoid exclusive repeated use of herbicides from the same herbicide group code. Alternate or tank mix with products from different herbicide group codes,
- Integrate the control methods (chemical, cultural, biological) into weed control programmes.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: Use only as directed.



FARM-AG



VULTURE 480 EC

Reg No/Nr: L10047 Act/Wet 36 of/van 1947

An emulsifiable concentrate of a systemic herbicide for the control of woody plants and weeds as indicated for forestry, grass pastures industrial areas and sugarcane.

'n Sistemiese emulgeerbare konsentraat onkruidodder vir die beheer van houtagtige plante en onkruid soos aangetoon in bosbou, grasweidings en nywerheidsgebiede.

HRAC HERBICIDE GROUP CODE	0	HRAC ONKRUIDODDERGROEPKODE
----------------------------------	----------	-----------------------------------

ACTIVE INGREDIENT:

triclopyr (pyridyloxy compound 480 g/ℓ triklopir (piridiloksieverbinding as butoxyzyl ethyl ester) as butoksie-etielester)

Manufacture Date:
Vervaardigingsdatum

Batch No.:
Lotnommer

NET VOLUME / NETTO VOLUME

SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE

REGISTERED & DISTRIBUTED BY / GEREISTREUR & VERSPREI DEUR:

Cedar Falls Properties 130 (Pty) Ltd

Co. Reg. No: 2006/031383/07

22 Burnside Drive, Old Mill Industrial Park, South Africa 4300 • Tel: 031 538 8700

UN No.: 3082



HARMFUL
SKADELIK



Use Precaution:

- Ensure that the person assigned to apply the product is properly trained in its use.

Compatibility:

- **VULTURE 480 EC** is compatible with Hyper Medium Oil (Reg. Nr.: L7355) and with formulations containing glyphosate.
- In case of any uncertainty contact the nearest **VULTURE 480 EC** supplier.

Mixing Instructions:

- For foliar applications, half-fill the spray tank with water, then add the required quantity of **VULTURE 480 EC** into the tank while stirring.
- While filling the spray tank with water to the required level, add Hyper Medium Oil at the rate indicated below.
- Maintain constant agitation.
- Ensure continuous agitation during filling and spraying operations.

Application Instructions:

- Avoid drift.
- Do not spray under conditions when spray drift to desirable vegetation may occur.
- Always calibrate the equipment under field conditions.

Foliar application:

(use only on actively growing plants with full leaf cover)

- Apply as a full cover spray to leaves and stems.
- Do not apply to wet foliage.
- Avoid spray drift onto non-target plants, other crops, grazing, rivers, dams or areas not under treatment.
- Do not apply if there is a probability rain shortly after application. Rain will have no effect on efficacy, provided the spray has dried on

the target plants.

- Use a knapsack sprayer with a solid cone nozzle e.g. Spraying Systems TG-1, Delavan CE1 or equivalent type, that will ensure an even coverage of the target area.
- Maintain a constant pressure of between 20 to 250 kPa.
- When applied using a tractor and boom, use about 350 lts per hectare or enough to give good foliar coverage.

Cut stump application:

- Apply within 12 hours of cutting/felling.
- When **VULTURE 480 EC** is applied in diesel it should not be applied to stumps that are wet from rain or dew.
- Apply the spray mixture at a low pressure of 10kPa and use solid cone nozzles e.g. Spraying Systems TG-1, Delavan or equivalent type.
- Apply the mixture to the point of run-off according to directions for stump application in the table below.

Basal Bark Treatment:

(for plants with a stem diameter of up to 10cm)

- **VULTURE 480 EC** in diesel should not be applied to stems if the bark is wet from rain or dew.
- Apply by paint brush or knapsack as a coarse low-pressure spray using narrow angle solid cone nozzle.
- Apply all round to the basal stem or trunk of the woody plant from soil level to a height indicated, as well as any exposed roots.
- In order to obtain satisfactory results a thorough wetting of the indicated area is imperative.
- Old rough bark may require more mixture than young or smooth bark.
- In the case of multi-stemmed plants, each stem should be treated separately.

DOSAGE RATES AND METHODS OF APPLICATION:**A. Foliar applications:**

Hyper Medium Oil should be added at a rate of 500 mℓ per 100 ℓ spray mixture for all foliar applications.

PLANT SPECIES	VULTURE 480 EC PER 100ℓ WATER	REMARKS
<i>Acacia cyclops</i> (Red eye)	500 mℓ (0,5 % mixture)	Only apply to young trees less than 2 metre in height.
<i>Acacia saligna</i> (Port Jackson willow)	600 mℓ (0.6% mixture)	
<i>Acacia dealbata</i> (Silver wattle)	250 – 750 mℓ (0,25 – 0,75% mixture)	Lower rate can be used on saplings less than 50 cm. Higher rate on young trees 0,5 to 2 metre tall.
<i>Acacia mearnsii</i> (Black wattle)		
<i>Asclepias fruticosa</i> (Shrubby milkweed)	375 mℓ (0,375 % mixture)	Apply to actively growing plants until the point of run-off.

PLANT SPECIES	VULTURE 480 EC PER 100ℓ WATER	REMARKS
<i>Caesalpinia decapatala</i> (Mauritius thorn)	500 mℓ (0,5 % mixture)	Apply to young, actively growing plants up to a maximum height of 3 metres. Taller plants must be cut back and regrowth treated only once it is 50 cm or higher.
<i>Chromolaena odorata</i> (Paraffin weed)	500 mℓ (0,5 % mixture)	
<i>Eucalyptus grandis</i> (Bluegum)	750 mℓ (0,75 % mixture)	Apply to leaves and stems when coppice is between 1,0 and 1,5 m in height. Ensure that surfaces are thoroughly wetted. Repeat treatment if regrowth occurs and reaches 0,4 metre.
<i>Pereskia aculeate</i> (Pereskia)	500 mℓ (0,5 % mixture)	Use a spray volume per hectare according to plant density. Plants should be reduced to a height of 2 m. All dead plant material should be removed and burned.
<i>Rubus cuneifolius</i> (American bramble)	500 mℓ (0,5 % mixture)	Treat young actively growing plants, using 300 to 400 litre spray mixture per treated hectare. Large plants should be slashed and treated when regrowth reaches 0,5 metre in height.
<i>Sesbania punicea</i> (Sesbania)	500 mℓ (0,5 % mixture)	Plants up to 2 metre high may be treated. Taller plants must be cut back and regrowth treated when at least 50 cm tall.
<i>Solanum mauritianum</i> (Bugweed)	500 mℓ (0,5 % mixture)	Treat plants up to 1,5 metre tall. Taller plants must be cut back and regrowth sprayed when at least 5 cm tall.
<i>Solanum sisymbriifolium</i> (Wild tomato)	500 mℓ (0,5 % mixture)	Apply to actively growing plants ensuring a thorough wetting. Limited regrowth may occur requiring a follow-up treatment.
<i>Stoebe plumose</i> (Slangbos)	500 mℓ (0,5 % mixture)	

B. Cut Stump Application:

PLANT SPECIES	VULTURE 480 EC PER 100ℓ DIESEL	REMARKS
<i>Acacia dealbata</i> (Silver wattle) <i>Acacia mearnsii</i> (Black wattle)	2,0 ℓ (2,0% mixture)	Cut 10 to 15 cm above ground level. Apply to cut surface and all exposed bark and root crown of stumps.
<i>Chromolaena odorata</i> (Trifid weed)	1,0 ℓ (1,0% mixture)	Cut stems approximately 10 cm above ground. Apply to all cut surfaces and all exposed bark and roots.
<i>Eucalyptus grandis</i> (Saligna gum)	2,0 ℓ (2,0 % mixture)	Treat the cleaned, active growing region (cambium) of the cut surfaces, all the bark, root crown and any exposed roots. Spray to the point of run-off. Monitor the area and re-apply to coppice if necessary.
<i>Prosopis spp</i> (Mesquite)	4,0 ℓ (4,0 % mixture)	Cut trees 10 to 15cm above ground. Apply to cut stump (bark, root crown and exposed roots). Monitor the area and repeat steps above to all regrowth.
<i>Sesbania punicea</i> (Sesbania)	0,5 ℓ (0.5 % mixture)	Cut stumps approximately 10 cm above ground level. Apply to cut stem surfaces and any exposed bark, rot crown and roots.
	OR 1,0 ℓ in water (1,0 % mixture)	For application in water only treat cut stem surfaces (not bark). Add wetting agent to mixture.

C. Basal Stem Applications:

PLANT SPECIES	VULTURE 480 EC PER 100ℓ DIESEL	REMARKS
<i>Acacia mearnsii</i> (Black wattle)	2,0 ℓ (2,0 % mixture)	Treat stems from approximately 25 cm downward. Also treat exposed root crown and roots ensuring thorough wetting of all plant parts treated. Deep rooted species may regrow. Foliar applications on all such regrowth is essential to ensure proper control.
<i>Cestrum laevigatum</i> (Inkberry)	1,0 ℓ (1,0 % mixture)	
<i>Melia azedarach</i> (Syringa)	2,0 ℓ (2,0 % mixture)	
<i>Robinia pseudoacacia</i> (Black locust)	2,0 ℓ (2,0 % mixture)	
<i>Schinus terebinthifolius</i> (Brazilian pepper tree)	2,0 ℓ (2,0 % mixture)	
<i>Solanum mauritianum</i> (Bugweed)	2,0 ℓ (2,0 % mixture)	

D. Sugarcane (NB: Do not use Hyper Medium Oil or any other wetter)

PLANT SPECIES	VULTURE 480 EC PER 100ℓ WATER	REMARKS
<i>Anredera basselloides</i> (Madeira vine)	500 mℓ	Apply to the weeds as a directed spot spray. Ensure thorough coverage of the target weeds.
<i>Physalis viscosa</i> (Sticky gooseberry)	(0.5% mixture)	

VULTURE 480 EC controls the trees and shrubs as listed in the tables above. The registration holder does not accept any responsibility for unlisted trees and shrub species not being controlled or failure of the product due to application methods used, not on this label or recommended by the registration holder.

NOTE: In situations where more than one species of target plant occurs contact your supplier or the registration holder. For assistance with calibration and spot treatment, contact your supplier or the registration holder.

WAARSKUWINGS:

- Skadelik indien ingesluk word.
- Konsentraat kan irritasie op vel en oë veroorsaak.
- **VLAMBAAR** – Moenie naby oop vure stoor nie.
- Stoor in 'n koel droë plek.
- Stoor ver weg van kos, voer, saad en ander landbouchemikalieë.
- Hou buite bereik van kinders, oningeligte persone en diere.
- Mag giftig wees vir visse.
- Herbetreding: Moenie behandelde area binne 1 dag na behandeling betree nie, behalwe wanneer beskermdes klere gedra word.
- **In geval van vergiftiging raadpleeg 'n geneesheer en stel hierdie etiket aan horn/haar beskikbaar.**

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid van toestande getoets is, waarborg die Registrasiehouer nie dat dit onder alle omstandighede doeltreffend sal wees nie, aangesien die uitwerking en resultaat daarvan beïnvloed kan word deur eksterne faktore soos abnormale grond, klimaats- en bergingstoestande, kwaliteit van water, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van die onkruid teen die betrokke middel, sowel as die metode, tyd en akkuraatheid van toediening. Verdermeer aanvaar die Registrasiehouer nie verantwoordelijkheid vir skade aan gewasse, plantegroei en die omgewing of skade teenoor mens en dier, of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim van die gebruiker om etiketaanwysings te volg of as gevolg van die ontstaan van toestande wat nie volgens die Registrasie voorsien kon word nie. Raadpleeg die verskaffer in die geval van enige onsekerheid.

VOORSORGMATREËLS:

Neem die volgende voorsorgmaatreëls, vir u eie veiligheid:

- Moenie eet, drink of rook terwyl die produk hanteer word nie.
- Moenie eet, drink of rook tydens die meng of toedien van die produk voordat hande en gesig gewas en skoon klere aangetrek is nie.
- Was hande en gesig na gebruik van die produk.
- Voorkom kontaminasie van kos, voer, drinkwater en eetgerei.

Tydens vermenging:

- Dra ondeurlaatbare handskoene en oogbeskerming.
- Was met seep en lopende water na toevallige velkontak.
- In geval van oogkontak, spoel oë onmiddellik met skoon lopende water, en verky mediese advies.

Tydens bespuiting:

- Moenie dampe of spuitnewel inasem nie.
- Vermy direkte kontak met die spuitmengsel.
- Dra geskikte beskermende klere.
- Verhoed spuitstofwegdrywing na ander gewasse, weidings, riviere, damme of areas wat nie behandel word nie.

Na bespuiting:

- Maak spuittoerusting deeglik skoon na gebruik en raak ontslae van spoelwater waar dit nie voedsel, ander gewasse, weiding, riviere of damme sal besoedel nie.

- Trek besoedelde klere uit en was dit.
- Was u hele liggaam, insluitende u hare en onder die vingernaels met seep en water.
- Spoel die lêë hours drie keer uit op die volgende manier: Keer die lêë houer om oor die spuit of mengten en laat toe om vir ongeveer 30 sekondes te dreineer nadat die vloei stadiger is en net druppels gesien kan word. Daarna moet die houer drie keer gespoel word met 'n volume water gelykstaande aan 'n minimum van 10% van die inhoud van die houer. Voeg die spoelwater by die inhoud van die spuitnek voordat die houer soos voorgeskryf vernietig word.
- Vernietig die lêë houer deur gate daarin te slaan en dit plat te slaan en raak ontslae daarvan op 'n veilige manier.
- Moet nooit die lêë houer hergebruik vir enige ander doeleindes nie.

SIMPTOME VAN MENSLIKE VERGIFTIGING:

Simptome wat verband hou met gebruik:

Velkontak: Irritasie op vel (jeukerigheid, rooiheid, pyn).

Oogkontak: Daar word nie verwag dat daar 'n merkbaar oogkontak gevaar sal voorkom onder verwagte toestande of normale gebruik nie.

Inname: Maagpyn, naarheid, probleme om asem te haal. Wanneer ingesluk word mag dit aanleiding gee tot longbeskadiging.

EERSTEHULP BEHANDELING:

Velkontak: Vir selfs geringe kontak, trek besoedelde klere, skoen en alle leergoedere uit. Was die vel saggies maar deeglik met water en 'n nie-skurende seep. Indien die irritasie van vel aanhou, moet mediese kundigheid ingewin word.

Oogkontak: Spoel die besoedelde oë onmiddellik af met sagvloeiende skoon water vir ongeveer 20 minute, terwyl die ooglede oopgehou word totdat daar geen meer bewyse van die chemiese stof oorby is. Indien irritasie nog voortduur, verky mediese hulp.

Inaseming: Verseker dat vars lug ingesam word. Verwyder die bron van besoedeling of beweeg die slagoffer na 'n plek waar daar vars lug is. Indien dit moeilik is om asem te haal, dien suurstof toe. Indien die persoon opgehou het om asem te haal, moet kunsmatige asemhaling toegepas word. Indien die asemhalingsprobleme voortduur, verky mediese hulp.

Inname: Moenie braking veroorsaak nie. Die slagoffer moet mond deeglik met water uitspoel. Verky mediese advies.

NOTA AAN GENESHEER:

Geen spesifieke teenmiddel is beskikbaar nie. Indien nodig moet behandeling simptomaties en ondersteunend toegepas word.

WEERSTANDSWAARSKUWING:

VULTURE 480 EC is 'n groepkode O onkruidodder. Enige onkruidbevolking mag individuele natuurlike weerstand teen **VULTURE 480 EC** en enige ander groepkode O onkruid toon. Die weerstandige individue kan uiteindelik die onkruidbevolking domineer indien hierdie onkruidodders herhaaldelik gebruik word. Hierdie weerstandige onkruid mag dalk nie deur **VULTURE 480 EC** of enige ander groepkode O onkruidodder beheer word nie.

Om onkruidodder weerstandigheid te verjaag:

- Vermoë uitsluitlike herhaalde gebruik van onkruidodders van dieselfde onkruid groepkode. Wissel af met of gebruik tenkmengsels van produkte in verskillende onkruidodder groepkodes.
- Integreer ander beheermaatreëls (chemise, verbouing en biologies) in onkruidodder programme.

Vir spesifieke inligting rakende weerstandsbestuur kan die Registrasiehouer van hierdie produk gekontak word.

GEBRUIKSAANWYSINGS: Gebruik slegs soos aangedui.

Wees Versigtig:

- Verseker dat die aangestelde persoon wat die produk moet aanwend behoorlik opgelei is om dit te gebruik.

Verenigbaarheid:

- **VULTURE 480 EC** is verenigbaar met Hiper Medium Olie (Reg. Nr.: L7355) en met formulerings met 'n gifosaat inhoud.
- In geval van enige onsekerheid moet die naaste **VULTURE 480 EC** verskaffer gekontak word.

Menginstruksies:

- Blaartoedienings, vul die spuittenk half-vol water en voeg die voorgeskrewe volume hoeveelheid **VULTURE 480 EC** by, terwyl dit geroer word.
- Terwyl die spuittenk met die voorgeskrewe hoeveelheid water gevul word, moet Hyper Medium Olie teen die aanbevole dosis hieronder bygevoeg word.
- Roer mengsel aanhoudend.
- Verseker dat die mengsel aanhoudend geroer word gedurende die volmaak en spuitoperasies.

Toedieningsinstruksies:

- Vermoë spuitnewel wegdrywing.
- Moenie spuit in ongunstige toestande waar spuitnewel kan wegdryf na nie-geteikende plante.
- Verseker dat die spuittoerusting altyd korrek gekalibreer word onder veldtoestande.

Blaartoedienings:

(Wend slegs aan op aktief groeiende plante wat volle drag blare het)

- Dien toe as 'n volle dekbespuiting op die blare en stamme.

- Moenie toedien terwyl die blare nat is nie.
- Verhoed spuitstofwegdrywing na gewasse, riviere, damme en gebiede wat nie behandel word nie.
- Moenie toedien indien daar 'n moontlikheid van reën kort na toediening is nie. Reën sal geen invloed op die effektiwiteit hê nie, mits die spuitstof droog is op die geteikende plante.
- Gebruik 'n rugsakspuit met 'n soliede spuitpunt bv. Spraying Systems TG-1, Delavan CE1 of soortgelyke spuitstukke, om te verseker dat die geteikende area eweredig gedek is.
- Handhaaf 'n konstante druk van 200 to 250 kPa.
- Tydens toediening moet 'n trekker en 'n spuitbalk gebruik word. Gebruik omtrent 350 l per hektaar of genoeg om goeie blaardekking te verskaf.

Gekapte stompbespuiting:

- Dien binne 12 ure na afkapping toe.
- Wanneer **VULTURE 480 EC** toegedien is met diesel behoort dit nie toegedien te word op nat stampe as gevolg van reën of dou nie.
- Dien die spuitmengsel toe teen 'n lae druk van 10kPa deur gebruik te maak van soliede keelspuitpunte bv. Spraying Systems TG-1, Delavan of soortgelyke spuitstukke.
- Dien toe tot die punt van afloop volgens die aanwysingstabel hieronder.

Basale Basbehandeling:

(vir plante met 'n stamdeursnee van 10cm)

- **VULTURE 480 EC** in diesel behoort nie toegedien te word op stamme wat nat is weens reën of dou nie.
- Dien toe met 'n verfkwas of knapsack as 'n growwe laedruk bespuiting.
- Dien toe rondom die basale stam of stomp vandie houtagtige plante, vanaf grondhoogte tot 'n hoogte soos aangedui, asook op enige ontblote wortels.
- Deeglike benutting van die aangeduide areas is noodsaaklik vir goeie beheer.
- Ou of growwe bas mag meer spuitstof benodig as jong of gladde bas.
- In die geval van meerstammige plante, behoort elke stam afsonderlik behandel te word.

TOEDIENINGSHOEVEELHEDE EN METODE VAN TOEDIENING:

A. Blaartoediening:

Hyper Medium Olie moet teen 'n dosis van 500 ml per 100 l spuitmengsel vir alle blaarbehandelings toegedien word.

PLANT SPESIES	VULTURE 480 EC PER 100l WATER	OPMERKINGS
Acacia cyclops (Rooikrans)	500 ml (0,5 % mengsel)	Behandel slegs jong bome minder as 2 meter in hoogte.
Acacia saligna (Port Jackson)	600 ml (0.6% mengsel)	

PLANT SPESIES	VULTURE 480 EC PER 100ℓ WATER	OPMERKINGS
Acacia dealbata (Silwer wattle)	250 – 750 mℓ (0,25 – 0,75% mengsel)	'n Laer dosis kan gebruik word op saailinge kleiner as 50 cm. Hoër dosisse op jong bome 0,5 tot 2 meter hoog.
Acacia mearnsii (Swart wattle)	375 mℓ (0,375 % mengsel)	Dien toe op aktief groeiende plante tot die punt van afloop.
Asclepias fruticosa (Tontelbossie)	500 mℓ (0,5 % mengsel)	Behandel jong bome tot maksimum 3 m hoog. Hoër plante moet teruggesny word en hergroei moet behandel word op 50 cm of hoër.
Caesalpinia decapatala (Mauritiusdoring)	500 mℓ (0,5 % mengsel)	
Chromolaena odorata (Paraffienbos)	500 mℓ (0,5 % mengsel)	
Eucalyptus grandis (Bloekom)	750 mℓ (0,75 % mengsel)	Dien toe op blare en stamme sodra hergroei tussen 1,0 en 1,5 m in hoogte is. Verseker dat die area deeglik benat word. Herhaal die toediening as hergroei weer voorkom en 0,4 meter bereik.
Pereskia aculeate (Pereskia)	500 mℓ (0,5 % mengsel)	Gebruik 'n spuitvolume per hektaar volgens die digtheid van die plante. Plante behoort verkort te word tot 'n hoogte van 2 m. Alle dooie plantmateriaal behoort verwyder of verbrand te word.
Rubus cuneifolius (Amerikaanse braam)	500 mℓ (0,5 % mengsel)	Behandel slegs jong aktief groeiende plante deur 300 to 400 liter spuitstof per behandelde hektaar te gebruik. Groter plante behoort afgekap te word en behandel te word sodra die hergroei 0,5 meter in hoogte bereik.
Sesbania punicea (Sesbania)	500 mℓ (0,5 % mengsel)	Plante met 'n hoogte tot 2 meter mag behandel word. Hoër plante moet teruggesny word en hergroei moet behandel word wanneer dit ten minste 50 cm hoog is.
Solanum mauritianum (Luisboom)	500 mℓ (0,5 % mengsel)	Behandel plant tot en met 1,5 meter hoog. Hoër plante moet teruggesny word en hergroei moet gespruit word wanneer dit ten minste 5 cm hoog is.
Solanum sisymbriifolium (Wilde tamatie)	500 mℓ (0,5 % mengsel)	
Stoebe plumose (Slangbos)	500 mℓ (0,5 % mengsel)	Dien toe tot aktief groeiende plante en verseker dat dit deeglik nat is. Beperkte hergroei mag voorkom wat 'n opvolg behandeling moet kry.

B. Gekapte Stompbehandeling:

PLANT SPESIES	VULTURE 480 EC PER 100ℓ DIESEL	OPMERKINGS
Acacia dealbata (Silwer wattle)	2,0 ℓ	Sny stampe af tot by 10 tot 15 cm bokant grondvlak. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels.
Acacia mearnsii (Swart wattle)	(2,0% mengsel)	
Chromolaena odorata (Paraffienbos)	1,0 ℓ (1,0% mengsel)	Sny stampe ongeveer 10 cm bokant die grond. Behandel die gesnyde oppervlakte asook alle bas, wortelkroon en blootgestelde wortels.
Eucalyptus grandis (Bloekom)	2,0 ℓ (2,0 % mengsel)	Behandel die skoon, aktief groeiende gedeelte (kambium) van die gekapte oppervlakte, al die bas, wortelkroon en enige blootgestelde wortels. Bespuit tot die punt van afloop. Hou die bespuitte area dop en opvolgbehandeling mag nodig wees.
Prosopis spp (Mesquite)	4,0 ℓ (4,0 % mengsel)	Sny bome 10 tot 15cm bokant die grond. Behandel dit gesnyde stomp (bas, wortelkroon en blootgestelde wortels). Hou die area dop en herhaal die stappe soos bo genoem in geval van hergroei.
Sesbania punicea (Sesbania)	0,5 ℓ (0,5 % mengsel)	Sny stampe ongeveer 10 cm bokant die grondvlak. Behandel gesnyde oppervlak en enige blootgestelde bas, wortelkroon en wortels.
	OF 1,0 ℓ in water (1,0 % mengsel)	Vir behandelings met water word slegs gesnyde oppervlakte behandel (nie die bas nie). Voeg water by mengsel.

C. Basale Stambehandelings:

PLANT SPESIES	VULTURE 480 EC PER 100ℓ DIESEL	OPMERKINGS
<i>Acacia mearnsii</i> (Swart wattle)	2,0 ℓ (2,0 % mengsel)	Behandel stamme van omtrent 25 cm afwaarts.
<i>Cestrum laevigatum</i> (Inkbessie)	1,0 ℓ (1,0 % mengsel)	Behandel ook blootgestelde wortelkroon
<i>Melia azedarach</i> (Sering)	2,0 ℓ (2,0 % mengsel)	en wortels en verseker deeglike benatting
<i>Robinia pseudoacacia</i> (Wit akasia)	2,0 ℓ (2,0 % mengsel)	van alle dele van die plant wat behandel
<i>Schinus terebinthifolius</i> (Brasiliaanse peperboom)	2,0 ℓ (2,0 % mengsel)	is. Diep gewortelde spesie mag hergroei.
<i>Solanum mauritianum</i> (Luisboom)	2,0 ℓ (2,0 % mengsel)	Blaarbehandelings op sulke hergroei is noodsaaklik en verseker behoorlike beheer.

D. Suikerriet (NB: Moenie Hyper Medium Olie of enige ander benatter gebruik nie)

PLANT SPESIES	VULTURE 480 EC PER 100ℓ WATER	OPMERKINGS
<i>Anredera basselloides</i> (Madeira vine)	500 mℓ	Dien toe op onkruid as 'n direkte kolspuit. Verseker deeglike bedekking van die geteikende onkruide.
<i>Physalis viscosa</i> (Klewerige appelliefie)	(0.5% mengsel)	

VULTURE 480 EC beheer die bome en struik soos gelys in die tabelle hierbo. Die Registrasiehouer aanvaar nie verantwoordelikheid vir ongelyste bome en struik spesies wat nie beheer is nie of waar die produk gefaal het danksy toedieningsmetodes gebruik wat nie op die etiket verskyn of aanbeveel word deur die Registrasiehouer nie.

NOTE: In situasies waar meer as een spesie van die geteikende plante voorkom, moet die verskaffer of Registrasiehouer gekontak word. Vir ondersteuning met kalibrasie en kolbehandeling, moet die verskaffer of Registrasiehouer gekontak word.



FARM-AG

VULTURE 480 EC

Reg No/Nr: L10047 Act/Wet 36 of/van 1947

An emulsifiable concentrate of a systemic herbicide for the control of woody plants and weeds as indicated for forestry, grass pastures industrial areas and sugarcane.

'n Sistemiese emulgeerbare konsentraat onkruidodder vir die beheer van houtagtige plante en onkruid soos aangetoon in bosbou, grasweidings en nywerheidsgebiede.

HRAC HERBICIDE GROUP CODE	0	HRAC ONKRUIDODDERGROEPKODE
----------------------------------	----------	-----------------------------------

ACTIVE INGREDIENT:

triclopyr (pyridyloxy compound 480 g/ℓ triklopir (piridiloksieverbinding as butoxyzyl ethyl ester) as butoksie-etielester)

Manufacture Date:
Vervaardigingsdatum

Batch No.:
Lotnommer

NET VOLUME / NETTO VOLUME

SEE ENCLOSED LEAFLET FOR FULL PARTICULARS
SIEN INGESLOTE VOUBILJET VIR VOLLE BESONDERHEDE

REGISTERED & DISTRIBUTED BY / GEREISTREUR & VERSPREI DEUR:

Cedar Falls Properties 130 (Pty) Ltd

Co. Reg. No: 2006/031383/07

22 Burnside Drive, Old Mill Industrial Park, South Africa 4300 • Tel: 031 538 8700

UN No.: 3082



HARMFUL
SKADELIK

